

Eumastós® gel

Breast gel

COMPOSITION

Each sachet of 2.5 g contains: Boswellia serrata, betaine, myo-inositol, Hamamelis virginiana, propylene glycol, hydroxyethyl cellulose, glycerin, ethoxydiglycol, potassium sorbate, sodium benzoate, EDTA, o-Cymen-5-ol, lactic acid, purified water.

CATEGORY

Single-use Medical Device.

PROPERTIES

EUMASTÓS® gel is a dermatological preparation in gel form whose local use helps maintain the normal physiological conditions of the mammary gland. The substances with antioxidant effect in EUMASTÓS® gel counteract free radicals and protect cell membranes by acting on the oxidative processes. EUMASTÓS® gel eases breast tenderness and pain for fast and long-lasting relief. EUMASTÓS® gel has a soothing effect on the skin area where it is applied.

PHARMACEUTICAL FORM

Gel.

CLINICAL INFORMATION

INDICATIONS

Resolution of pain caused by surgical wounds or diagnostic procedures. Treatment and prevention of premenstrual and/or cyclical mastalgia (breast pain). Soothes breast tenderness and disorders related to non-cancerous breast diseases. Adult population.

CONTRA-INDICATIONS

Known hypersensitivity to the components.

SIDE-EFFECTS

No side-effects have been reported to date. Tell your doctor and pharmacist about any undesirable effects you suspect may have been caused by using the product.

INCOMPATIBILITIES / INTERACTIONS

To date, there are no known interactions with other devices, substances or medicinal products.

EFFECTS ON THE ABILITY TO DRIVE AND USE MACHINES

No effects on the ability to drive and use machines have been reported.

DOSAGE AND METHOD OF ADMINISTRATION

Unless advised otherwise by your doctor, apply the contents of one gel sachet once a day on each breast, massaging in gently until completely absorbed. For cyclical mastalgia (breast pain), apply from day 10 to day 25 of the menstrual cycle.



WARNINGS

For single use. For external use. Do not use the product if you are allergic to any of the components. The use of any product for topical use, especially if prolonged, can lead to sensitisation; if this occurs, discontinue treatment and speak to your doctor to start suitable therapy. No injuries from overdose have been reported under normal conditions of use. Do not use the product when pregnant or breastfeeding. Do not use the product after the expiry date shown on the package. Do not use the product if the packaging is not intact. Keep out of reach of children. Do not swallow.

PRECAUTIONS FOR STORAGE

Store the product at room temperature ($\leq 25\text{ }^{\circ}\text{C}$), in a cool, dry place protected from light and away from heat.

PACKAGE

30 sachet of 2.5 g.

DISPOSAL

There are no special requirements to be followed for disposal of the product. Check local regulations in force.



Leaflet paper PAP 22

Recycling - Please check the regulations in your municipality



Lo.Li. Pharma S.r.l.

Via Sabatino Gianni, 14 - 00156 Rome - Italy

